arXiv:astro-ph/0105040v2 3 May 2001

Ionization Structure and the Reverse Shock in E0102-72

K.A. Flanagan, C.R. Canizares, D.S. Davis, D. Dewey, J.C. Houck,
M.L. Schattenburg

Center for Space Research, Massachusetts Institute of Technology
Cambridge, MA 02139

Abstract. The young oxygen-rich supernova remnant E0102-72 in the Small Magellanic Cloud has been observed with the High Energy Transmission Grating Spectrometer of Chandra. The high resolution X-ray spectrum reveals images of the remnant in the light of individual emission lines of oxygen, neon, magnesium and silicon. The peak emission region for hydrogen-like ions lies at larger radial distance from the SNR center than the corresponding helium-like ions, suggesting passage of the ejecta through the "reverse shock". We examine models which test this interpretation, and we discuss the implications.

BACKGROUND

1E0102.2-7219 (hereinafter referred to as E0102-72) is a young (∼1000 years) oxygen-rich supernova remnant (SNR) in the Small Magellanic Cloud. E0102-72 has been observed at all wavelengths: in the optical [1] [2], UV [3] [4] , radio [5], and in the X-ray [6] [7] [8] [9] [10] [11] [12] [13]. Gaetz *et al.* have looked at direct Chandra images of E0102-72 and noted the radial variation with energy bands centered on OVII and OVIII, suggesting an ionizing shock propagating inward. This paper addresses this issue through analysis of the high resolution dispersed spectrum, and reinforces this interpretation.

THE HIGH RESOLUTION X-RAY SPECTRUM

Shown in Figure 1 is the dispersed high resolution X-ray spectrum of E0102-72. The Chandra observation was made using the High Energy Transmission Gratings (HETG) in conjunction with the Advanced CCD Imaging Spectrometer (ACIS-S). The HETG separates the X-rays into their distinct emission lines, forming separate images of the remnant with the X-ray light of each line. The dispersed spectrum contains lines of highly ionized oxygen, neon, magnesium and silicon (not shown in the figure). Detailed study of the line images shows they are distorted along the dispersion axis due to Doppler shifts associated with high velocity material. Details and an overview of results are given in Canizares *et al.* elsewhere in these proceedings. Here we examine the apparent differences in ring radii seen in Figure 1. As discussed in detail below, these are likely caused by the changing ionization state, suggesting the passage of the supernova ejecta through the reverse shock.



FIGURE 1. A portion of the high resolution spectrum formed by the medium energy gratings. At right in the figure is the zeroth order, which combines all energies in an undispersed image.

THE IONIZATION/SHOCK STRUCTURE OF E0102-72

Figure 2 compares helium-like and hydrogen-like lines of oxygen. The dispersed OVII Resonance line from the helium-like triplet (left) is displayed next to the hydrogen-like OVIII Ly α line (right) on the same spatial scale. The ring diameter of the hydrogen-like line is obviously larger than that of the helium-like line. The ring diameters of these lines were measured by tracing the distribution in the direction orthogonal to the dispersion axis, as shown in Figure 3. (By taking the distribution in the cross-dispersion direction, the result is largely independent of Doppler shift.) The He-like OVII distribution is nestled cleanly within the OVIII, and is measurably narrower in Figure 3. The ring diameters for all the bright lines were measured



FIGURE 2. The OVII Resonance line is emitted from a region of smaller radius than that of OVIII Lyman α.



FIGURE 3. The OVII Resonance distribution is narrower than that of OVIII Lyman α.

similarly, and the results are shown on Figure 4. *All of the hydrogen-like lines lie outside the corresponding helium-like lines.* After passage of a shock, an ionizing plasma at a fixed electron temperature T_e will achieve a He-like state before it reaches a H-like state. This is shown in Figure 5, where the appropriate timescale is the ionization parameter $\tau = n_e t$, where t is the time since passage of the shock and n_e is the electron density [14]. Since the H-like state lies outward of the He-like state in E0102-72, this suggests the action of a shock moving *inward* relative to the ejecta - the 'reverse shock' which is the standard model for the mechanism which heats SNR ejecta to X-ray temperatures.



FIGURE 4. Ring diameter vs energy for the bright lines. Note that all the hydrogen-like lines (connected by the top curve) lie outside their helium-like counterparts.



FIGURE 5. Helium-like NeIX Resonance line reaches its peak emissivity at logτ=10.0, earlier than hydrogen-like NeX Lyman α, which peaks at about log τ=11.0 sec/cm^3



FIGURE 6. Measured diameter vs τ at peak emissivity. O7 is an upper limit for τ.

We applied a simple model in which a homogenous mix of ejecta is shocked to a fixed T_e of about 1 keV or log $T_e = 10^{7.05}$ (as suggested by our global NEI analysis [11]). We estimated τ by finding the peak emissivity (as in Figure 5). A plot of measured diameters as a function of τ is shown in Figure 6. Although any workable model must consider many parameters, the monotonic behavior suggests that these differences in ring diameter are attributable to the ionization structure resulting from the reverse shock.

WORK IN PROGRESS AND FUTURE WORK

Our current model calls for further work. For example, by taking the ratio of the histograms for O8 Lyman α and O7 Resonance from Figure 3, we can examine the allowed region of paramater space defined by T_e and τ *as a function of radial position.* (This approach as it pertains to the global spectrum is described by Davis *et al* [11] elsewhere in these proceedings.) By considering a likely density gradient in the ejecta and obtaining an appropriate estimate for electron density, we might obtain a corresponding estimate for the reverse shock velocity. We plan to apply this shock velocity information to see how it fits with the spatial extent of the bright ejecta and the presumed age of the remnant.

Acknowledgements We thank Glenn Allen, Norbert Schulz and Sara-Anne Taylor for helpful discussions. We are grateful to the CXC group at MIT for their advice and assistance. This work was prepared under NASA contract NAS8-38249 and SAO SV1-61010.

REFERENCES

1. Dopita, M.A., Tuohy, I.R. & Mathewson, D.S., 1981, *ApJ*, **248**, L105 (1981)
2. Tuohy, I.R. & Dopita, M.A.,*ApJ*,**268**, L11 (1983)
3. Blair, W.P., Raymond, J.C., Danziger, J. & Matteucci, F., *ApJ*, **338**, 812 (1989)
4. Blair, W.P., Morse, J.A., Raymond, J.C., Kirshner, R.P., Hughes, J.P., Dopita, M.A., Sutherland, R.S., Long, K.S. & Winkler, P.F., *ApJ*, **537**, 667 (2000)
5. Amy, S.W. & Ball, L., *ApJ*, **411**, 761 (1993)
6. Seward, F.D. & Mitchel, M., *ApJ*, **243**, 736 (1981)
7. Hayashi, I., Koyama, K., Masanobu, O., Miyata, E., Tsunemi, H., Hughes, J.P. & Petre, R., *PASJ*, **46**, L121 (1994)
8. Gaetz, T.J., Butt, Y.M., Edgar, R.J., Eriksen, K.A., Plucinsky, P.P., Schlegel, E.M. & Smith, R.K., *ApJ*, **534**, L47 (2000)
9. Hughes, J.P. 1988, in *Supernova Remnants and the Interstellar Medium*, ed. R.S. Roger & T.L. Landecker Cambridge: Cambridge Univ. Press, p. 125
10. Hughes, J.P., Rakowski, C.E. & Decourchelle, A. *ApJ*, (2000) in press
11. Davis, D.S., Flanagan, K.A., Houck, J.C., Canizares, C.R, Allen, G.E., Schulz, N.S., Dewey, D. & Schattenburg, M.L., *these proceedings.*
12. Canizares, C.R, Flanagan, K.A., Davis, D.S., Dewey, D., & Houck, J.C., *these proceedings.*
13. Rasmussen, A. *et al.*, *these proceedings.*
14. Hughes, J.P. & Helfand, D.J., *ApJ*, **291**, 544 (1985)